



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### **Differential effects of soil waterlogging on herbaceous and woody plant communities in a Neotropical savanna**

**Citation for published version:**

De Oliveira Xavier, R, Leite, MB, Dexter, K & Da Silva Matos, DM 2019, 'Differential effects of soil waterlogging on herbaceous and woody plant communities in a Neotropical savanna', *Oecologia*, pp. 1-13.  
<https://doi.org/10.1007/s00442-019-04423-y>

**Digital Object Identifier (DOI):**

[10.1007/s00442-019-04423-y](https://doi.org/10.1007/s00442-019-04423-y)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Oecologia

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



Differential effects of soil waterlogging on herbaceous and woody plant communities in a Neotropical savanna

Rafael de Oliveira Xavier<sup>1,2\*</sup>, Marcelo Boccia Leite<sup>3</sup>, Kyle Dexter<sup>2,4</sup>, Dalva Maria da Silva Matos<sup>4</sup>

1- Department of Ecology, Biosciences Institute, University of São Paulo, Rua do matão 123, São Paulo, SP, 05508-090, Brazil.

2- School of Geosciences. The University of Edinburgh, Edinburgh EH9 3JN, UK

3- Ecology and Conservation Lab, Department of Hydrobiology, Federal University of São Carlos, Washington Luís highway, km 235, São Carlos, SP 13565-905, Brazil.

4- Royal Botanic Garden Edinburgh, 20a Inverleith row, EH3 5LR, Edinburgh, UK

\*Corresponding author: [rxavier@ib.usp.br](mailto:rxavier@ib.usp.br), phone: +55 11 979854169

R.O. Xavier, M. B. Leite and D.M. Silva-Matos conceived the ideas and designed the methodology; R.O. Xavier and M. B. Leite collected the data; R. O. Xavier and K. Dexter analysed the data; R. O. Xavier and K. Dexter led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication

## ABSTRACT

The impacts of soil properties and fire regime on Neotropical savannas are well-known, but the importance of hydrological regime for plant species assembly has received less attention. Here we assessed changes in diversity patterns of herbaceous and woody communities along a water table gradient in a fire-excluded Neotropical savanna. We found that increased waterlogging of soils was associated with declines in both herbaceous and woody species richness. Woody species richness decreased once the water table depth is less than 4 m and no woody species occurred once water table depth was less than 23 cm. Herbaceous communities remained species rich until the shallowest water table depth, where there is flooding at some point in the year, and even there, over a dozen species occurred. Woody species that occurred in areas with shallower water tables were a nested subset of those in areas with deeper water tables. In contrast, herbaceous communities showed turnover over the hydrological gradient, with distinct species specialised for different water table levels. However, we found that those specialists are restricted to few evolutionary lineages, evidenced by increased phylogenetic clustering over the water table gradient in herbaceous communities. We suggest that evolutionarily conserved hydrological niches define the herbaceous layer over the hydrological gradient, whereas only generalist woody species persist under high water tables. Our findings show that the effect of soil waterlogging differs between the herbaceous and woody layer of savannas, indicating that these communities will respond differently to shifts in the hydrological regime under future environmental change.

Keywords: water table depth, hydrological niches, turnover, phylogenetic.

## Introduction

The varying patterns of plant community assembly across environmental gradients have inspired important ecological concepts and have been a recurring subject of research since the earliest ecological studies (Clements 1936; Keddy 1992; Laliberté et al. 2014; Whittaker 1956), and there is increasing interest in understanding plant community assembly in light of ongoing global environmental changes (Lewthwaite et al. 2017). The effects of larger-scale geographic gradients in water availability on community composition and species richness have been demonstrated in a wide range of ecosystems, from tropical forests (Esquivel-Muelbert et al. 2017; Pyke et al. 2001) and seasonally dry ecosystems (Scholes et al. 2002) to riparian forests (Garssen et al. 2015) and floodplains (Casanova and Brock 2000; McGinness et al. 2018; Stevenson et al. 2018). In addition, local variation in hydrological regime can shape plant community composition at individual sites (Araya et al. 2011; Kirkman et al. 2001; Silvertown et al. 1999), which has led researchers to define distinct hydrological niches for different species (Silvertown et al. 2015). However, research in this area has been limited by the difficulty of quantifying hydrological regime. Although assessing soil texture, a proxy for how quickly water percolates through the soil (Wakindiki and Ben-Hur 2002), and topography, which generally relates to the water table depth (Desbarats et al. 2002), is relatively easy, actually measuring water table depth over annual cycles requires substantially more time and financial investment (Morgan and Stolt 2004). Spatial and temporal variation in water tables has been quantified in temperate plant communities, where it is consistently found to have strong effects on plant community composition (Allen-Diaz 1991; Dwire et al. 2006; Shafroth et al. 2000; Silvertown et al. 1999). Similar studies are scarce in the tropics (although see Moulatlet et al. 2014; Targhetta et al. 2015), despite the tropics housing the overwhelming majority of plant diversity.

Tropical savannas occur in seasonally dry climates, and typically show a continuous herbaceous layer and a discontinuous woody canopy (Sarmiento 1984). Factors driving the occurrence of savanna *versus* forest have been extensively investigated (de Castro and Kauffman 1998; Sankaran et al. 2005), and recent large-scale studies have highlighted the role of fire regime and herbivory (Hempson et al. 2015; Lehmann et al. 2014; Staver et al. 2011). However, seasonal flooding and soil waterlogging, potentially in conjunction with fire occurrence, are also related to the occurrence of savannas in large areas of South America, including the *llanos* of Venezuela and Colombia (Sarmiento and Pinillos 2001), the Rupupuni savannas of Guyana (Eden 1970) and the Llanos de Moxos of Bolivia (Hamilton et al. 2004). There are also extensive areas of hydrologically determined savannas in Brazil (Batalha et al. 2005; Cianciaruso et al. 2005; Tannus and Assis 2004), yet most studies of the woody cover and composition of Brazilian savanna have focused on fire regime as a driving variable (Cianciaruso et al. 2012; Moreira 2000; Silva and Batalha 2010).

The few studies of hydrological gradients in Brazilian savannas (e.g. Villalobos-Vega et al. 2014, Leite et al 2018) have focused on the woody plant layer, while much of plant species richness of Brazilian savannas lies in the herbaceous layer (Rodrigues Munhoz and Felfili 2007; Tannus and Assis 2004). In fact, herbaceous species can account for over 90% of the plant species richness (Mendonça et al. 1998; Rezende et al. 2008) and most of the primary productivity (Lloyd et al. 2008). In other ecosystems, diversity patterns of herbaceous and woody species are differentially related to environmental variation (Moro et al. 2015; Yan et al. 2013). This is especially expected under distinct hydrological regimes, considering the recurrent differences in water acquisition strategies between herbaceous and woody species (Dodd et al. 1998; Moro et al. 2015; Walter 1979).

In this study, we aimed to assess the effect of water table depth and other abiotic drivers on patterns of species richness and lineage diversity in both woody and herbaceous plant communities along a hydrological gradient in fire-excluded savannas in Southern Brazil, where shallow water tables are thought to be a key driver of open vegetation (Leite et al. 2018). Although both recent fire suppression and past fire regimes are expected to affect current plant community patterns in these savannas, as with any other savanna (Bond and Parr 2010; Brockway and Lewis 1997), the overall study site has experienced little changes in woody cover during the last three decades despite marked variation in woody cover within the site. In other words, the spatial pattern of variation in woody cover has remained relatively static, suggesting that the hydrological regime may be the major force affecting vegetation structure at the site (Leite et al. 2018). Therefore, our study site offers a rare opportunity of investigating the effect of a hydrological gradient on community diversity patterns while controlling for potential additional effects of fire regime.

We specifically aimed to answer the following questions: i) Is the water table depth a major driver of variation in the species richness of plant communities?; ii) Is there turnover in plant species composition along water depth gradients, or are the plant species in more species-poor communities a nested subset of those in more species-rich communities?; iii) Are the plant communities in areas with high water tables comprised of species from few or many evolutionary lineages, i.e. do these plant communities show phylogenetic clustering? iv) Do these patterns differ for herbaceous and woody plant communities? Flooding gradients often lead to high species and phylogenetic turnover in plant communities (Garssen et al. 2015; Silvertown et al. 1999; Tanentzap and Lee 2016), and hence we expect large differences in species and phylogenetic community structure between sites with contrasting hydrological regimes. In addition,

recent research at our study site has demonstrated that the density of the woody layer decreases as water tables become shallower (Leite et al. 2018), so we also expect matching decreases in woody plant species richness. In contrast, we hypothesize that changes in diversity patterns at the herbaceous layer will be present, if at all, along a smaller extent of the water depth gradient, considering that poorly drained Neotropical grasslands and savannas are often species-rich ecosystems (Bueno et al. 2014; Munhoz and Felfili 2006; Killeen and Hinz 1992, Tannus and Assis 2004).

## **Material and methods**

### *Study site*

This study was carried out at the Itirapina Ecological Station (IES), a 2,300 ha protected area in southeast Brazil (Fig. 1a). The local climate is Cwa (Köppen) and the average annual rainfall and temperature during the study period were 1,613 mm and 20.6 °C, respectively (IFSP 2018). There is typically a favourable water balance during the rainy season and a short period of water deficit in the dry season (Xavier et al. 2017). The prevalent soil is an entisol with at least 90% sand (Leite et al. 2018). The IES contains a gradient in vegetation physiognomy that is typical of broader savanna vegetation in Brazil, from grasslands with few shrubs and trees (open savanna - *campo sujo*) to a more closed savanna with higher tree density (dense savanna - *cerrado sensu stricto*) and including an intermediate type (savanna - *campo cerrado*) (Coutinho 1978). The IES maintains some of the last remnants of tropical grasslands in southeast Brazil, where open savannas have been affected by degradation and woody encroachment (Durigan et al. 2007; Stevens et al. 2016). There are no detailed data on the disturbance regime before the IES was established (early 1980s), although the lack of past fire management policies suggest that fire used to occur at the reserve prior to this period.

Managers have excluded fire from the IES during the last 25 years. Aerial and satellite imagery taken since the mid-1980s suggest that the distribution of open savannas and grasslands in the IES has changed little during the last 30 years (Leite et al. 2018).

We selected 19 sites along a water table depth gradient (Fig. 1a). Vegetation along this gradient varies from wet and marshy grasslands with no woody species to savannas with variable woody cover (campo sujo, campo cerrado and cerrado *sensu stricto* – *sensu* Coutinho (1978), Electronic Supplemental Material ESM1). Although we initially aimed to sample an equal the number of sites within each of these three savanna categories, we found that woody cover varies widely within each savanna category (Leite et al. 2018). Likewise, we found that grassland types initially categorized based on visual differences in their flooding regime were in effect distributed along a hydrological gradient, including distinct annual periods of soil waterlogging and standing water (Xavier et al. 2017). Considering these limitations on identifying *a priori* physiognomies or vegetation types at our study site and our aim of assessing community responses to water table depth, our study focus on assessing diversity patterns of woody and plant communities occurring along a continuum of woody cover and hydrological regimes. Sites initially classified into the same vegetation type were always at least 500 m from one another. None of the sites have experienced fire during the last two decades.

At each site, we identified and counted all woody individuals with basal diameter greater than 3 cm in six randomly allocated 10 x 10 m plots (Fig. 1b). In order to sample the herbaceous community, in each of these plots we also established six regularly distributed 1 x 1 m subplots spaced 2 m apart, where we identified and counted all individuals of herbaceous species (Fig. 1b, Electronic Supplemental Material ESM2). We assumed that all the stems connected aboveground or at the soil



level belonged to the same individual; turfs of caespitose grasses and sedges with no aboveground connections between them were considered distinct individuals. Individuals were identified to the species level in the field or with further support of previous surveys in the study site, plant collections and consultation with taxonomists (Tannus and Assis 2004). We followed the nomenclature of The Plant List (2017). Unidentified species were classified into morphospecies within genera or families. Across all sites, we could not identify 828 herbaceous individuals to the family level (3.8 % of all herbaceous individuals sampled). Although these individuals were not included in the analysis, most of these morphotypes were single occurrences and hence their absence is unlikely to strongly affect inference on patterns of community assembly at our study sites. Based on these identifications and field observations, species were classified as trees, shrubs or herbs; subshrubs and woody species with no permanent aboveground woody stem were included in the herbaceous layer plant community as they interact primarily with the graminoid and forb species that are typically dominant at this layer. Specimens were deposited in the herbarium of the Federal University of São Carlos, Brazil.

#### *Phylogenetic and Environmental Data*

We created a phylogeny that included all species across the plots using the online tool Phylomatic (Webb and Donoghue 2005), based on the Angiosperm Phylogeny Group's APGIII consensus tree (R20120829). Node ages were estimated using the *bladj* algorithm in the software Phylocom 4.2. (Webb et al. 2008) with age constraints from Gastauer et al. (2017). This phylogeny was then pruned to a woody plant species phylogeny and an herbaceous plant species phylogeny, in order to generate separate null expectations for woody and herbaceous plant communities when quantifying phylogenetic community structure.

During the peak of the dry season in 2011 (August), we used a manual auger to drill one 40 mm-wide well down to the groundwater just outside five of the six 10 x 10 m vegetation sampling plots at each site (Electronic Supplemental Material ESM2). At sites where initial drilling showed the water table to be deeper than the maximum drilling capacity of our equipment (7 m), we drilled wells in only two of the six surveyed 10 x 10 m plots. We measured the groundwater depth every two weeks from October 2011 to October 2013 using a Solinst Water Level Meter 102 500' measuring tape. We used the minimum water table depth over this period as an explanatory variable because it represents the maximum waterlogging or highest water table that plants at a given site experience (Xavier et al. 2016). We averaged values across the different plots at a given site. The maximum capacity of the drilling equipment (7.5 m) was used as the minimum water table depth estimate for the sites where the groundwater was too deep to be measured.

We collected 15 soil samples from 0 to 30 cm depth at each study site, which were thoroughly mixed to produce a composite sample by site (see Electronic Supplemental Material ESM3 for the abiotic characterization of each site). Soil texture (sand, silt and clay content) and soil chemical composition (P, N, K, Mg, Al, Ca, pH and organic matter) were obtained following Camargo et al. (2013).

#### *Data analysis*

Our main response variables in analyses were the species richness, species diversity and lineage diversity of herbaceous and woody plant communities. Given an equal sample effort in terms of area sampled, our species richness values can be directly compared across sites. To quantify species diversity, we used the Simpson Index ( $1 - D$ ), which is not only a measure of species richness but also reflects how evenly

individuals are distributed across species. Phylogenetic diversity, or the total branch length of a phylogeny comprising the species at a given site, is the most basic measure of lineage diversity; however, this measure is strongly correlated with species richness, and high variation in species richness across communities, as we have at our sites, can obscure any patterns in variation of lineage diversity. We therefore used a null model to calculate the standardized effect size of phylogenetic diversity, based on reshuffling the tips of the phylogeny 1000 times. We refer to this standardized effect size as lineage diversity (c.f. Honorio Coronado et al. 2015; Rezende et al. 2017). It is also a measure of phylogenetic community structure, with negative values reflecting phylogenetic clustering and positive values reflecting phylogenetic overdispersion (Webb et al. 2002). We did not use alternative measures of phylogenetic community structure, such as the standardised effect size of the mean phylogenetic distance between species in a community (also referred to as the net relatedness index), because these alternative measures can be highly sensitive to the distribution of species among the three major clades of angiosperms (magnoliids, monocots and eudicots; see Honorio Coronado et al. 2015) and our herbaceous and woody plant communities differ substantially in the distribution of their constituent species across these clades. This would therefore complicate comparison of results for woody and herbaceous plant communities. Phylogenetic diversity and its effect size were calculated separately for tree and herbaceous species based on pruned trees comprising only tree and herbaceous species, respectively. These calculations were performed in the R package “picante” (Kembel et al. 2010).

Many of our soil variables covary strongly with each other and we therefore used biological reasoning to select a subset of the most pertinent explanatory variables. For example, the abundances of individual base cations are positively correlated with

each other, so we used the sum of bases as an explanatory variable in analyses. Also, the soil cation exchange capacity, base saturation, organic matter content and pH were strongly correlated with each other (see Electronic Supplementary Material ESM3 for soil properties from each study site). We included the soil base saturation as an explanatory variable as the variable most likely to be mechanistically related to plant community composition. The minimum water table depth and certain soil properties (N, sum of bases, Al) showed strongly right-skewed distributions in preliminary analyses and were therefore log-transformed prior to linear modelling to reduce heteroscedasticity.

We modelled the effect of water table depth and soil properties on species richness, species diversity and lineage diversity using generalized linear models with a negative binomial distribution for species richness (as it represents count data) and a Gaussian distribution for species and lineage diversity. Models were constructed separately for woody and herbaceous plant communities. Explanatory variables were scaled to have the same mean and variance to allow direct comparison of their regression coefficients as a measure of their relative effect on community composition. Given that our overarching aim was to assess the effect of water regime on community composition, we compared the Bayesian Information Criterion (BIC) of models including only water table depth as an explanatory variable with models including water table depth and each of the other soil properties in turn. Limited sample size in terms of study sites ( $N = 19$ ) prevented us from considering more complex models with more than two explanatory variables, or interactions. However, our approach does serve to test if our results for water table depth are robust to including the major axes of soil variation as covariates. When presenting results, we focus on presenting the best-fitting model for water table depth and one additional soil variable. Semivariograms showed

no clear spatial patterns in raw variables or residuals. Further, accounting for spatial autocorrelation did not improve model estimation in generalized linear models (following Zuur et al. 2009), and spatial autocorrelation was therefore not included in the final models presented here.

In order to visualise in greater detail how species richness, species diversity and lineage diversity vary with the original non log-transformed water table depth values, we used additive models with thin plate regression splines as smoothing functions with the R package “mgcv” (Wood 2006), without defining a maximum number of basis dimensions. We used this approach because additive models assume no *a priori* relationship between explanatory and response variables (Zuur et al. 2009). Therefore, they are a flexible alternative for identifying ecological thresholds and different types of biologically meaningful non-linear responses (Francesco Ficetola and Denoël 2009; Zuur et al. 2009), which are often found along extensive hydrological gradients such as at our study site (Dwire et al. 2004; Lite and Stromberg 2005).

We followed the approach of Leibold and Mikkelsen (2002) to assess if there is community turnover or nestedness along the water table gradient, separately for herbaceous and woody plant communities. We sorted the community matrix according to increasing water table depth and calculated the number of times a species was replaced by another between sites with adjacent water table depth; the resulting average turnover value was then compared to the values obtained from 1000 random communities created by permuting the original matrix (Leibold and Mikkelsen 2002). Significant differences between these values indicate that the community follows a turnover pattern, whereas similar values suggest a nested pattern. We performed these analyses using the function “Turnover” in the R package “metacom” (Dallas 2014). We also quantified the amount of nestedness in woody and herbaceous plant using the

NODF metric, which is based on decreasing fill and paired overlap in the community matrix (Almeida-Neto et al. 2008). Values obtained for each layer were compared to null expectations based on 30000 random communities that preserved the number of species occurrences and row and column totals (Ulrich et al. 2009). These analyses were performed in the R packages “vegan” (Oksanen et al. 2013) and “bipartite” (Dormann et al. 2008).

Herbaceous and woody morphotypes that could not be identified to the genus or family level were included in nestedness analyses. Matrices for the herbaceous layer included the whole water table gradient, whereas for the woody layer, they only included sites where woody species occurred (11 out of 19 sites). All analyses were performed in the R statistical environment (R\_Core\_Team 2017), with further support of packages cowplot (Wilke 2016) and ggplot2 (Wickham 2010) for graphic presentation of results.

## **Results**

We identified, at least to the family level, 22,652 individuals from 50 families and 210 species and morphospecies, including 154 herbs and 66 trees. The most common families in terms of abundance and species richness were Myrtaceae, Asteraceae and Fabaceae in the woody layer, and Poaceae and Cyperaceae in the herbaceous layer (see Electronic Supplemental Material ESM3 for species composition at each study site).

Species richness and species diversity were strongly correlated and similarly related to environmental variables, and we therefore only present results for species richness here (see Electronic Supplemental Material ESM4 for results based on species diversity). Species richness and lineage diversity of the herbaceous layer were closely

related to water table depth. Most of the best models in terms of BIC values also included soil properties, but the effect of these variables was always smaller than that showed by water table depth and did not affect the sign or significance of the water table depth effect (Table 1). Tree species richness was also strongly related to water table depth and weakly related to soil base saturation, whereas lineage diversity of woody plants was largely unrelated to water table depth or soil factors we measured (Table 1).

Generalised additive models (GAMs) showed that the diversity patterns of herbaceous and woody layer had different thresholds of response to water table depth (Fig. 2, Table 2). Although the herbaceous communities on sites subjected to flooding (i.e. negative water table depth) exhibited much lower species richness (points in bottom left of Fig. 2a), additive models only captured a weak linear relationship between water table and species richness. The failure to statistically detect this clearly non-linear relationship is likely related to the very small difference in the measured water table depth values between species-rich grasslands with very shallow water tables and species-poor seasonally flooded grasslands (Fig. 2a). Additive models showed that the lineage diversity of the herbaceous layer increased sharply as water table depth increased until approximately 1 m and remained stable thereafter (Fig. 2c). Woody species richness increased as the water table depth increased to about 4 m, and then showed no clear association with water table depth (Fig. 2b). Woody plant lineage diversity showed no clear relationships with water table depth (Fig. 2d).

The community structure of the herbaceous and woody layer also showed different patterns along the water table gradient (Fig. 3). Although certain generalist herb species occurred from grasslands with shallow water table to savannas with deep water table (Fig. 3a), the number of species replacements in the herbaceous layer

between sites with adjacent water table depth was significantly higher than expected according to a null model ( $z=-4.76$   $p<0.001$ ), suggesting a turnover pattern. In contrast, the number of replacements in the woody layer did not differ from expected according to a null model ( $z=0.64$ ,  $p=0.520$ ), suggesting a nested pattern along the water table gradient (Fig. 3b). Consistent with these findings, the amount of nestedness in the herbaceous layer (NODF=31.10) was significantly lower than expected based on a null model ( $z=-4.45$ ,  $p=0.003$ ) and considerably higher than in the woody layer (NODF=49.50), although the amount of nestedness at the woody layer did not differ from the null expectation ( $z=0.53$ ,  $p=0.678$ ).

## Discussion

Our study shows that seasonally high water tables, which lead to soil waterlogging, have major impacts on both herbaceous and woody plant communities, although the nature of these impacts differs. As shown in previous research at this site (Leite et al. 2018), woody plant species richness declines as water tables become shallower, with no woody plants able to persist as adult trees ( $> 3$  cm diameter at ground level) in areas where the water table comes within 20 cm of the ground surface (see also Fig. 2b). In contrast, many herbaceous species (more than a dozen) can persist at all water table levels, although herbaceous species richness does halve at the shallowest water table depths, where seasonal flooding occurs (Fig. 2a). The herbaceous species that persist at the shallowest water table depths are distinct from those that occur on higher and drier ground (Fig. 3a), and analyses of phylogenetic clustering show that these species come from relatively few evolutionary lineages. In contrast, woody species show a nested pattern of species distributions over hydrological gradients (Fig. 3b), and species occurring at shallower water tables do not seem restricted to particular evolutionary lineages.



348        As we expected, the richness at the woody layer was severely limited by shallow  
349        water table, suggesting intolerance to seasonal waterlogging. The flooding regime often  
350        has large effects on riparian woody plant communities (da Paz et al. ; Garssen et al.  
351        2015; Garssen et al. 2017), and recent studies have shown a negative effect of shallow  
352        water table on the tree species richness of savannas (Villalobos-Vega et al. 2014). We  
353        found no trees on sites where the minimum annual water table level was very close to or  
354        above the soil surface (i.e. where flooding occurred). These results are somewhat  
355        surprising, because in South America many seasonally flooded savannas have been  
356        described (Batalha et al. 2005; Bueno et al. 2014; Sarmiento et al. 2004), although these  
357        studies did not measure water table depth directly. Considering the negative effects of  
358        flooding on plant survival and performance, which require specific adaptations  
359        (Mommer et al. 2006), we hypothesize that extended time periods of shallow water  
360        tables and flooding are strong abiotic filters that, at least in some areas, cannot be  
361        overcome by woody species. Clearly, most tree species present in the species pool for  
362        our study area cannot survive such challenging hydrological conditions. We did find  
363        that a few tree species could occur where the water table remained a few centimeters  
364        deeper than the soil surface, as has been found for other poorly drained Neotropical  
365        savannas (Batalha et al. 2005; Bueno et al. 2014). Once water tables were deeper than 3  
366        m, we found that minimum water table depth had little effect on tree species richness.  
367        Shallow water tables may be buffering open savannas in our study site against woody  
368        encroachment associated with fire exclusion, which has been occurring in well-drained  
369        Neotropical savannas (Durigan and Ratter 2016; Stevens et al. 2016). However, our  
370        findings also show that small changes in the water table dynamics of sites with shallow

water table could have large effects on the species richness of woody plant communities in hydrologically-determined savannas.

Contrasting with the effects on woody species richness, there was no association between water table depth and the lineage diversity of woody communities. Although we found that certain tree species and families (e.g. Myrtaceae, Bignoniaceae, Fabaceae) only occurred where the water table was deeper than 3 m, a few species from multiple lineages (e.g. *Tibouchina stenocarpa*, *Lippia velutina*, *Moquiniastrum polymorphum*) occurred in savannas with both shallow and deep water tables. This is consistent with the complex interaction of anatomical, morphological and physiological adaptations associated with flood-tolerance (Garssen et al. 2015; Kolb and Joly 2009; Kozłowski 1997), which tend not to be phylogenetically clustered (Mommer et al. 2006). Although it is possible that this flooding tolerance generally occurs at taxonomic levels lower than the resolution in our phylogeny (family and genus-level resolution), phylogenies at the same resolution have been used to associate environmental variation and phylogenetic composition (Fine and Kembel 2011; Kubota et al. 2014). In addition, a study based on a fully-resolved phylogeny also found no evidence of phylogenetic clustering along a soil moisture gradient, suggesting that responses to distinct water regimes may be evolutionarily labile in woody plant lineages (Araya et al. 2012). Although assessing if this is a widespread pattern in tropical savannas subjected to high water tables would require investigations at larger spatial scales, our findings suggest that future changes in the hydrological regime in our study site would affect the woody community lineage diversity less than the species richness.

We found that woody species that do occur in areas with high water tables are simply a subset of species from higher and drier areas that have the necessary physiological adaptations to manage this stressful, water-logged environment. Previous

studies also found little turnover of tropical woody species along water availability gradients (Esquivel-Muelbert et al. 2016). In effect, we found that poorly drained savannas were dominated by *Moquiniastrum polymorpha* (Less.) G. Sancho, which also occurs in well-drained areas. This species shows high plasticity in leaf structure, which has been associated with its occurrence in both Neotropical savannas and forests (Rossatto and Kolb 2010). Similarly, certain woody species from the family Melastomataceae, which were common where water tables were high in our study area, are common in Neotropical palm swamps and show leaf anatomy plasticity what seem to favour their occurrence in poorly drained sites (Somavilla and Graciano-Ribeiro 2011). Although plasticity may not be associated with environmental tolerance (Dostál et al. 2016), plasticity on anatomical, physiological and morphological traits, also called low oxygen escape syndrome, allows species without metabolic adaptations to flooding to persist in flood-prone environments (Bailey-Serres and Voesenek 2008). For example, *Sinojackia huangmeiensis* J.W. Ge & X.H. Yao showed distinct seed morphology and composition when subjected to soil waterlogging (Wei et al. 2018). Therefore, phenotypic plasticity, rather than evolutionarily rare adaptations to soil waterlogging, may enable the persistence of these savanna trees species under shallow water tables.

#### *Herbaceous communities*

Our data suggest that herbaceous species richness was only reduced over the shallowest water tables, where seasonal flooding occurred. This is expected considering the large negative effect on plant performance and survival of severe soil anoxia and hypoxia associated with flooding, whereas more herbaceous species are expected to occur where there is only seasonal soil waterlogging (Wegner 2010). These findings are consistent with the typically high herbaceous species richness found in other poorly drained Brazilian savannas (Bueno et al. 2014; Munhoz and Felfili 2006; Rodrigues

Munhoz and Felfili 2007; Tannus and Assis 2004). Previous studies in temperate grasslands found similar diversity patterns (Dwire et al. 2006), suggesting that seasonally shallow water tables may not be a constraint on herbaceous species richness in a wide range of ecosystems.

In contrast to effects on species richness, the lineage diversity of the herbaceous layer was reduced under high water tables. Consistent with this effect, the herbaceous layer of savannas with water tables deeper than 3 m included families (e.g. Menispermaceae, Bromeliaceae) absent from sites with shallow water tables. In Neotropical savannas, many perennial herbs and subshrubs are found with relatively large underground structures that support both water uptake and rapid post-fire regeneration (de Castro and Kauffman 1998; Sarmiento 1984). These species may be unable to overcome the hypoxic conditions in sites subjected to soil waterlogging (Bailey-Serres and Voesenek 2008; Pezeshki 2001; Wegner 2010). In our study, these lineages seem to be replaced by a few lineages of small perennial and annual herbs on sites with intermediate water table depth (e.g. Onagraceae, Xyridaceae, Eriocaulaceae), which maintained a high species richness; this likely led to the only slightly lower lineage diversity at these sites compared to the better drained savannas. In contrast, continuous waterlogging and periodic flooding require specific mechanisms to overcome severe soil hypoxia (Bailey-Serres and Voesenek 2008; Kozłowski 1984), so that lineage diversity decreases markedly towards more severe flooding (Tanentzap and Lee 2016). Consistent with this expectation, a few graminoid species and taxa typical from ephemeral wetlands (e.g. *Hydrocotyle*, *Maiacaceae*) (Deil 2005) occurred in sites subjected to flooding.

Differently from the woody layer, we found compositional turnover in the herbaceous layer over the hydrological gradient, possibly because herbaceous and

woody plants are differentially affected by high water table. Poor drainage is expected to be a constraint to many cerrado woody species, which often have a deep underground system (Oliveira et al. 2005). Conversely, herbaceous species are often shallow rooted (Rawitscher 1948) and are only expected to be affected by soil waterlogging where the water table is right below the soil surface. In seasonally dry ecosystems, high water tables may even favour the occurrence of herbaceous species with poor tolerance to water deficit, as it may increase the moisture at the superficial soil layer by capillary action (Moore 1939). In effect, there is no annual period of water deficit in most of our study sites subjected to seasonal waterlogging (Xavier et al. 2017). Therefore, we hypothesize that conditions associated with high water tables support specific herbaceous species, and competitive ability is expected to be important for their persistence (Grime 1988). In summary, herbaceous species are most likely to occur where the hydrological regime enables them to perform close to their ecological optima (Silvertown et al. 1999), leading to low niche overlap and species segregation along water table depth gradients (Silvertown et al. 1999). The long-term fire suppression at our study sites has possibly contributed to the emergence of these patterns, as it is expected to favour the most competitive species under different hydrological conditions, to the detriment of generalist and disturbance-prone herbaceous species (Grime 1988; Tilman 1985). Regardless of this possible local effect, our findings highlight the functionally unique composition, and high conservation value, of Neotropical grasslands with high water tables.

We conclude that water table depth is the major abiotic factor affecting savanna plant communities at our study site, but this effect varies between the herbaceous and woody layer. There were no trees under hydrological flooding, and only a few woody species, albeit from multiple lineages, persisted at high water tables. Indeed, shallow

water table is a major constraint to the occurrence of tree species in savannas, which may be buffering open savannas in our study site against woody encroachment associated with fire exclusion. Conversely, only actual flooding limited the richness of herbaceous plant communities, whilst high water tables excluded certain lineages and favoured others. For herbaceous plant communities, there was species turnover along the water table depth gradient, consistent with species having distinct hydrological niches. Although further studies should investigate if these diversity patterns are widespread in savanna communities that occur in poorly-drained sites, and how they interact with fire regime, these findings clearly suggest that herbaceous and woody plant communities will respond differently to shifting hydrological regimes under global environmental change.

#### Acknowledgements

We are thankful to the Coordination for the Improvement of Higher Education Personnel (CAPES) for the funding granted to RO Xavier and MB Leite and to the State of São Paulo Research Foundation (FAPESP) for the scholarship granted to RO Xavier (Grant 2011/21019-6). KG Dexter was supported by a Leverhulme International Academic Fellowship during the time this work was completed. We also thank the São Paulo Forestry Institute for the research permit and to the Itirapina Ecological Station staff for logistical assistance, and the Royal Botanical Garden Edinburgh for providing full access to its collection. We thank A. Magurran for her initial contribution on the conceptual direction of the study manuscript. J. Veldman and an anonymous reviewer made valuable suggestions to improve later versions of the manuscript.

#### Disclosure of potential conflicts of interest

Authors declare no conflict of interest.

## References

- Allen-Diaz BH (1991) Water table and plant species relationships in Sierra Nevada meadows. *American Midland Naturalist* 126:30-43
- Almeida-Neto M, Guimarães P, Guimarães PR, Loyola RD, Ulrich W (2008) A consistent metric for nestedness analysis in ecological systems: reconciling concept and measurement. *Oikos* 117:1227-1239. doi: 10.1111/j.0030-1299.2008.16644.x
- Araya YN, Silvertown J, Gowing DJ, McConway KJ, Linder HP, Midgley G (2012) Do niche-structured plant communities exhibit phylogenetic conservatism? A test case in an endemic clade. *Journal of Ecology* 100:1434-1439.
- Araya YN, Silvertown J, Gowing DJ, McConway KJ, Linder HP, Midgley G (2011) A fundamental, eco-hydrological basis for niche segregation in plant communities. *New Phytologist* 189:253-258
- Bailey-Serres J, Voisenek LACJ (2008) Flooding Stress: Acclimations and Genetic Diversity. *Annual Review of Plant Biology* 59:313-339. doi: 10.1146/annurev.arplant.59.032607.092752
- Batalha MA, Cianciaruso MV, Silva IA, Delitti WBC (2005) Hyperseasonal cerrado, a new brazilian vegetation form. *Brazilian Journal of Biology* 65:735-738
- Bond WJ, Parr CL (2010) Beyond the forest edge: Ecology, diversity and conservation of the grassy biomes. *Biological Conservation* 143:2395-2404. doi: <https://doi.org/10.1016/j.biocon.2009.12.012>
- Brockway DG, Lewis CE (1997) Long-term effects of dormant-season prescribed fire on plant community diversity, structure and productivity in a longleaf pine wiregrass ecosystem. *Forest Ecology and Management* 96:167-183. doi: [https://doi.org/10.1016/S0378-1127\(96\)03939-4](https://doi.org/10.1016/S0378-1127(96)03939-4)

Bueno ML, Damasceno-Junior GA, Pott A, Pontara V, Seleme EP, Fava WS, Salomão  
 AK, Ratter JA (2014) Structure of arboreal and herbaceous strata in a neotropical  
 seasonally flooded monodominant savanna of *Tabebuia aurea*. *Brazilian Journal of*  
*Biology* 74:325-337

Casanova MT, Brock MA (2000) How do depth, duration and frequency of flooding  
 influence the establishment of wetland plant communities? *Plant Ecology*: 147:237-  
 250

Cienciaruso MV, Antônio Batalha M, Aurélio da Silva I (2005) Seasonal variation of a  
 hyperseasonal cerrado in Emas National Park, central Brazil. *Flora - Morphology,*  
*Distribution, Functional Ecology of Plants* 200:345-353. doi:  
<http://dx.doi.org/10.1016/j.flora.2005.02.001>

Cienciaruso MV, Silva IA, Batalha MA, Gaston KJ, Petchey OL (2012) The influence  
 of fire on phylogenetic and functional structure of woody savannas: Moving from  
 species to individuals. *Perspectives in Plant Ecology, Evolution and Systematics*  
 14:205-216. doi: <https://doi.org/10.1016/j.ppees.2011.11.004>

Clements FE (1936) Nature and structure of the climax. *Journal of ecology* 24:252-284

Coutinho LM (1978) O conceito do cerrado. *Revista Brasileira de Botânica* 1:17-23

da Paz AA, Ribeiro C, Azevedo AA, de lima ER, da Silva Carmo FM Induced flooding  
 as environmental filter for riparian tree species. *Environmental and Experimental*  
*Botany* 139: 31-38

Dallas T (2014) metacom: an R package for the analysis of metacommunity structure.  
*Ecography* 37:402-405

de Castro EA, Kauffman JB (1998) Ecosystem structure in the Brazilian Cerrado: a  
 vegetation gradient of aboveground biomass, root mass and consumption by fire.  
*Journal of Tropical Ecology* 14:263-283. doi: doi:null



- Deil U (2005) A review on habitats, plant traits and vegetation of ephemeral wetlands - a global perspective. *Phytocoenologia* 35:533-706. doi: 10.1127/0340-269X/2005/0035-0533
- Desbarats AJ, Logan CE, Hinton MJ, Sharpe DR (2002) On the kriging of water table elevations using collateral information from a digital elevation model. *Journal of Hydrology* 255:25-38. doi: [https://doi.org/10.1016/S0022-1694\(01\)00504-2](https://doi.org/10.1016/S0022-1694(01)00504-2)
- Dodd MB, Lauenroth WK, Welker JM (1998) Differential water resource use by herbaceous and woody plant life-forms in a shortgrass steppe community. *Oecologia* 117:504-512. doi: 10.1007/s004420050686
- Dormann CF, Gruber B, Fründ J (2008) Introducing the bipartite package: analysing ecological networks. *R news* 8:8-11
- Dostál P, Fischer M, Chytrý M, Prati D (2016) No evidence for larger leaf trait plasticity in ecological generalists compared to specialists. *Journal of Biogeography*:n/a-n/a. doi: 10.1111/jbi.12881
- Durigan G, Siqueira MFd, Franco GADC (2007) Threats to the Cerrado remnants of the state of São Paulo, Brazil. *Scientia Agricola* 64:355-363
- Dwire KA, Kauffman JB, Baham JE (2006) Plant species distribution in relation to water-table depth and soil redox potential in montane riparian meadows. *Wetlands* 26:131-146
- Eden M (1970) Savanna vegetation in the northern Rupununi, Guyana. *Journal of Tropical Geography* 30:17-28
- Esquivel-Muelbert A, Baker TR, Dexter KG, Lewis SL, ter Steege H, Lopez-Gonzalez G, Monteagudo Mendoza A, Brien R, Feldpausch TR, Pitman N, Alonso A (2016) Seasonal drought limits tree species across the Neotropics. *Ecography* 40:618-629. doi: 10.1111/ecog.01904

571 Fine PVA, Kembel SW (2011) Phylogenetic community structure and phylogenetic  
 572 turnover across space and edaphic gradients in western Amazonian tree  
 573 communities. *Ecography* 34:552-565. doi: 10.1111/j.1600-0587.2010.06548.x  
 574 Francesco Ficetola G, Denoël M (2009) Ecological thresholds: an assessment of  
 575 methods to identify abrupt changes in species–habitat relationships. *Ecography*  
 576 32:1075-1084. doi: 10.1111/j.1600-0587.2009.05571.x  
 577 Garssen AG, Baattrup-Pedersen A, Voosenek LACJ, Verhoeven JTA, Soons MB (2015)  
 578 Riparian plant community responses to increased flooding: a meta-analysis. *Global*  
 579 *Change Biology* 21:2881-2890. doi: 10.1111/gcb.12921  
 580 Garssen AG, Baattrup-Pedersen A, Riis T, Raven BM, Hoffman CC, Verhoeven JT,  
 581 Soons MB (2017) Effects of increased flooding on riparian vegetation: Field  
 582 experiments simulating climate change along five European lowland streams.  
 583 *Global change biology* 23:3052-3063  
 584 Grime JP (1988) The CSR model of primary plant strategies—origins, implications and  
 585 tests. *Plant evolutionary biology*. Springer, pp 371-393  
 586 Hamilton SK, Sippel SJ, Melack JM (2004) Seasonal inundation patterns in two large  
 587 savanna floodplains of South America: the Llanos de Moxos (Bolivia) and the  
 588 Llanos del Orinoco (Venezuela and Colombia). *Hydrological Processes* 18:2103-  
 589 2116. doi: 10.1002/hyp.5559  
 590 Hempson GP, Archibald S, Bond WJ (2015) A continent-wide assessment of the form  
 591 and intensity of large mammal herbivory in Africa. *Science* 350:1056-1061  
 592 Honorio Coronado EN, Dexter KG, Pennington RT, Chave J, Lewis SL, Alexiades MN,  
 593 Alvarez E, Alves de Oliveira A, Amaral IL, Araujo-Murakami A, Arets EJ (2015)  
 594 Phylogenetic diversity of Amazonian tree communities. *Diversity and Distributions*  
 595 21:1295-1307

596 IFSP (2018) Dados meteorológicos - Estação Ecológica de Itirapina. Instituto Florestal  
 597 de São Paulo, [http://www.iflorestal.sp.gov.br/Itirapina/dados\\_meteorologicos.html](http://www.iflorestal.sp.gov.br/Itirapina/dados_meteorologicos.html)  
 598 Keddy PA (1992) Assembly and response rules: two goals for predictive community  
 599 ecology. *Journal of Vegetation Science* 3:157-164  
 600 Kembel SW, Cowan PD, Helmus MR, Cornwell WK, Morlon H, Ackerly DD,  
 601 Blomberg SP, Webb CO (2010) Picante: R tools for integrating phylogenies and  
 602 ecology. *Bioinformatics* 26:1463-1464. doi: 10.1093/bioinformatics/btq166  
 603 Kirkman LK, Mitchell RJ, Helton RC, Drew MB (2001) Productivity and species  
 604 richness across an environmental gradient in a fire-dependent ecosystem. *American*  
 605 *Journal of Botany* 88:2119-2128  
 606 Kolb RM, Joly CA (2009) Flooding tolerance of *Tabebuia cassinoides*: Metabolic,  
 607 morphological and growth responses. *Flora - Morphology, Distribution, Functional*  
 608 *Ecology of Plants* 204:528-535. doi: <http://dx.doi.org/10.1016/j.flora.2008.07.004>  
 609 Kozłowski T (1984) Plant responses to flooding of soil. *BioScience* 34:162-167  
 610 Kozłowski T (1997) Responses of woody plants to flooding and salinity. *Tree*  
 611 *physiology* 17:490  
 612 Kubota Y, Hirao T, Fujii S-j, Shiono T, Kusumoto B (2014) Beta diversity of woody  
 613 plants in the Japanese archipelago: the roles of geohistorical and ecological  
 614 processes. *Journal of Biogeography* 41:1267-1276. doi: 10.1111/jbi.12290  
 615 Laliberté E, Zemanik G, Turner BL (2014) Environmental filtering explains variation in  
 616 plant diversity along resource gradients. *Science* 345:1602-1605  
 617 Lehmann CE, Anderson TM, Sankaran M, Higgins SI, Archibald S, Hoffmann WA,  
 618 Hanan NP, Williams RJ, Fensham RJ, Felfili J, Hutley LB (2014) Savanna  
 619 Vegetation-Fire-Climate Relationships Differ Among Continents. *Science* 343:548-  
 620 552. doi: 10.1126/science.1247355

- Leibold MA, Mikkelsen GM (2002) Coherence, species turnover, and boundary clumping: elements of meta-community structure. *Oikos* 97:237-250. doi: doi:10.1034/j.1600-0706.2002.970210.x
- Leite MB, Xavier RO, Oliveira PTS, Silva FKG, Matos DMS (2018) Groundwater depth as a constraint on the woody cover in a Neotropical Savanna. *Plant and Soil*:1-15
- Lewthwaite JMM, Debinski DM, Kerr JT (2017) High community turnover and dispersal limitation relative to rapid climate change. *Global Ecology and Biogeography* 26:459-471. doi: 10.1111/geb.12553
- Lite SJ, Stromberg JC (2005) Surface water and ground-water thresholds for maintaining *Populus*–*Salix* forests, San Pedro River, Arizona. *Biological conservation* 125:153-167
- Lloyd J, Bird MI, Vellen L, Miranda AC, Veenendaal EM, Djagbletey G, Miranda HS, Cook G, Farquhar GD (2008) Contributions of woody and herbaceous vegetation to tropical savanna ecosystem productivity: a quasi-global estimate†. *Tree Physiology* 28:451-468. doi: 10.1093/treephys/28.3.451
- McGinness HM, Arthur AD, Davies M (2018) Flood regimes driving vegetation and bird community transitions in semiarid floodplain woodlands. *Ecohydrology* 11:e1954. doi: doi:10.1002/eco.1954
- Mendonça RC, Felfili JM, Walter BM, Silva M, Rezende A, Filgueiras TS, Nogueira PE (1998) Flora vascular do cerrado. *Cerrado: ambiente e flora*. Embrapa-Cpac Planaltina
- Mommer L, Lenssen JPM, Huber H, Visser EJW, Kroon HD (2006) Ecophysiological determinants of plant performance under flooding: a comparative study of seven

plant families. *Journal of Ecology* 94:1117-1129. doi: doi:10.1111/j.1365-2745.2006.01175.x

Moore R (1939) Water conduction from shallow water tables. *Hilgardia* 12:383-426

Moreira AG (2000) Effects of fire protection on savanna structure in Central Brazil. *Journal of Biogeography* 27:1021-1029. doi: 10.1046/j.1365-2699.2000.00422.x

Morgan C, Stolt M (2004) A comparison of several approaches to monitor water-table fluctuations. *Soil Science Society of America Journal* 68:562-566

Moro MF, Silva IA, Araújo FSd, Nic Lughadha E, Meagher TR, Martins FR (2015) The Role of Edaphic Environment and Climate in Structuring Phylogenetic Pattern in Seasonally Dry Tropical Plant Communities. *PLOS ONE* 10:e0119166. doi: 10.1371/journal.pone.0119166

Moulatlet GM, Costa FRC, Rennó CD, Emilio T, Schietti J (2014) Local Hydrological Conditions Explain Floristic Composition in Lowland Amazonian Forests. *Biotropica* 46:395-403. doi: doi:10.1111/btp.12117

Munhoz C, Felfili J (2006) Floristics of the herbaceous and subshrub layer of a moist grassland in the Cerrado biosphere reserve (Alto Paraíso de Goiás), Brazil. *Edinburgh journal of botany* 63:343-354

Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MH, Wagner H (2013) *vegan: Community Ecology Package*. R package version 2.0-10.

Oliveira RS, Bezerra L, Davidson EA, Pinto F, Klink CA, Nepstad DC, Moreira A (2005) Deep root function in soil water dynamics in cerrado savannas of central Brazil. *Functional Ecology* 19:574-581. doi: 10.1111/j.1365-2435.2005.01003.x

668 Pezeshki SR (2001) Wetland plant responses to soil flooding. *Environmental and*  
 669 *Experimental Botany* 46:299-312. doi: [https://doi.org/10.1016/S0098-](https://doi.org/10.1016/S0098-8472(01)00107-1)  
 670 8472(01)00107-1  
 671 Pyke CR, Condit R, Aguilar S, Lao S (2001) Floristic composition across a climatic  
 672 gradient in a neotropical lowland forest. *Journal of Vegetation Science* 12:553-566.  
 673 doi: 10.2307/3237007  
 674 R\_Core\_Team (2017) R: A language and environment for statistical computing. R  
 675 Foundation for Statistical Computing. Vienna, Austria  
 676 Rawitscher F (1948) The Water Economy of the Vegetation of the 'Campos Cerrados'  
 677 in Southern Brazil. *Journal of Ecology* 36:237-268. doi: 10.2307/2256669  
 678 Mendonça RD, Felfili JM, Walter BM, Silva-Júnior MD, Rezende AV, Filgueiras TD,  
 679 Nogueira PE, Fagg CW (2008) Flora vascular do bioma cerrado. *Cerrado: ecologia*  
 680 *e flora*. Embrapa, Brasília:1028-1059  
 681 Rezende VL, Dexter KG, Pennington RT, Oliveira-Filho AT (2017) Geographical  
 682 variation in the evolutionary diversity of tree communities across southern South  
 683 America. *Journal of Biogeography* 44:2365-2375. doi: doi:10.1111/jbi.13013  
 684 Rodrigues Munhoz CB, Felfili JM (2007) Florística do estrato herbáceo-subarbustivo de  
 685 um campo limpo úmido em Brasília, Brasil. *Biota Neotropica* 7  
 686 Rossatto DR, Kolb RM (2010) *Gochnatia polymorpha* (Less.) Cabrera (Asteraceae)  
 687 changes in leaf structure due to differences in light and edaphic conditions. *Acta*  
 688 *Botanica Brasilica* 24:605-612  
 689 Sankaran M, Hanan NP, Scholes RJ, Ratnam J, Augustine DJ, Cade BS, Gignoux J,  
 690 Higgins SI, Le Roux X, Ludwig F, Ardo J (2005) Determinants of woody cover in  
 691 African savannas. *Nature* 438:846

- Sarmiento G (1984) *The Ecology of Neotropical Savannas*. Harvard University Press, Cambridge
- Sarmiento G, Pinillos M (2001) Patterns and processes in a seasonally flooded tropical plain: the Apure Llanos, Venezuela. *Journal of Biogeography* 28:985-996. doi: 10.1046/j.1365-2699.2001.00601.x
- Sarmiento G, Pinillos M, Silva MP, Acevedo D (2004) Effects of soil water regime and grazing on vegetation diversity and production in a hyperseasonal savanna in the Apure Llanos, Venezuela. *Journal of Tropical Ecology* 20:209-220
- Scholes R, Dowty P, Caylor K, Parsons DAB, Frost PGH, Shugart H (2002) Trends in savanna structure and composition along an aridity gradient in the Kalahari. *Journal of Vegetation Science* 13:419-428
- Shafroth PB, Stromberg JC, Patten DT (2000) Woody riparian vegetation response to different alluvial water table regimes. *Western North American Naturalist*:66-76
- Silva IA, Batalha MA (2010) Phylogenetic structure of Brazilian savannas under different fire regimes. *Journal of Vegetation Science* 21:1003-1013. doi: doi:10.1111/j.1654-1103.2010.01208.x
- Silvertown J, Araya Y, Gowing D, Cornwell W (2015) Hydrological niches in terrestrial plant communities: a review. *Journal of Ecology* 103:93-108. doi: doi:10.1111/1365-2745.12332
- Silvertown J, Dodd ME, Gowing DJG, Mountford JO (1999) Hydrologically defined niches reveal a basis for species richness in plant communities. *Nature* 400:61. doi: 10.1038/21877
- Somavilla NS, Graciano-Ribeiro D (2011) Análise comparativa da anatomia foliar de Melastomataceae em ambiente de vereda e cerrado sensu stricto. *Acta Botanica Brasilica* 25:764-775

- Staver AC, Archibald S, Levin SA (2011) The Global Extent and Determinants of Savanna and Forest as Alternative Biome States. *Science* 334:230-232. doi: 10.1126/science.1210465
- Stevens N, Lehmann CER, Murphy BP, Durigan G (2016) Savanna woody encroachment is widespread across three continents. *Global Change Biology*:n/a-n/a. doi: 10.1111/gcb.13409
- Stevenson PR, Aldana AM, Cárdenas S, Negret PJ (2018) Flooding and soil composition determine beta diversity of lowland forests in Northern South America. *Biotropica* 50:568-577. doi: doi:10.1111/btp.12541
- Tanentzap AJ, Lee WG (2016) Evolutionary conservatism explains increasing relatedness of plant communities along a flooding gradient. *New Phytologist*:n/a-n/a. doi: 10.1111/nph.14167
- Tannus JLS, Assis MA (2004) Composição de espécies vasculares de campo sujo e campo úmido em área de cerrado, Itirapina - SP, Brasil. *Brazilian Journal of Botany* 27:489-506
- Targhetta N, Kesselmeier J, Wittmann F (2015) Effects of the hydroedaphic gradient on tree species composition and aboveground wood biomass of oligotrophic forest ecosystems in the central Amazon basin. *Folia Geobotanica* 50:185-205. doi: 10.1007/s12224-015-9225-9
- Tilman D (1985) The Resource-Ratio Hypothesis of Plant Succession. *The American Naturalist* 125:827-852. doi: 10.1086/284382
- Ulrich W, Almeida-Neto M, Gotelli NJ (2009) A consumer's guide to nestedness analysis. *Oikos* 118:3-17. doi: 10.1111/j.1600-0706.2008.17053.x
- Villalobos-Vega R, Salazar A, Miralles-Wilhelm F, Haridasan M, Franco AC, Goldstein G (2014) Do groundwater dynamics drive spatial patterns of tree density and



diversity in Neotropical savannas? *Journal of Vegetation Science* 25:1465-1473.  
doi: 10.1111/jvs.12194

Wakindiki I, Ben-Hur M (2002) Soil mineralogy and texture effects on crust  
micromorphology, infiltration, and erosion. *Soil Science Society of America Journal*  
66:897-905

Walter H (1979) *Vegetation of the Earth and Ecological Systems of the Geo-Biosphere.*,  
Second Edition edn. Springer Verlag, Berlin-Heidelberg, New York

Webb CO, Ackerly DD, Kembel SW (2008) Phylocom: software for the analysis of  
phylogenetic community structure and trait evolution. *Bioinformatics* 24:2098-2100

Webb CO, Ackerly DD, McPeck MA, Donoghue MJ (2002) Phylogenies and  
community ecology. *Annual Review of Ecology and Systematics* 33:475-505

Webb CO, Donoghue MJ (2005) Phylomatic: tree assembly for applied phylogenetics.  
*Molecular Ecology Resources* 5:181-183

Wegner LH (2010) Oxygen transport in waterlogged plants. In: Mancuso SC, S. (ed)  
*Waterlogging Signalling and tolerance in Plants*. Springer, Berlin

Wei X, Liu M, Wang S, Jiang M (2018) Seed morphological traits and seed element  
concentrations of an endangered tree species displayed contrasting responses to  
waterlogging induced by extreme precipitation. *Flora* 246-247:19-25. doi:  
<https://doi.org/10.1016/j.flora.2018.07.001>

Whittaker RH (1956) *Vegetation of the Great Smoky Mountains*. Ecological  
Monographs 26:1-80. doi: 10.2307/1943577

Wickham H (2010) ggplot2: elegant graphics for data analysis. *J Stat Softw* 35:65-88

Wilke CO (2016) cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2'.  
R package version 0.7. 0

Wood S (2006) *Generalized Additive Models: An Introduction with R*. CRC Press

- Xavier RdO, Leite MB, da Silva-Matos DM (2016) Stress responses of native and exotic grasses in a Neotropical savanna predict impacts of global change on invasion spread. *Austral Ecology*:n/a-n/a. doi: 10.1111/aec.12475
- Xavier RO, Leite MB, da Silva-Matos DM (2017) Stress responses of native and exotic grasses in a Neotropical savanna predict impacts of global change on invasion spread. *Austral Ecology* 42:562-576
- Yan Y, Yang X, Tang Z (2013) Patterns of species diversity and phylogenetic structure of vascular plants on the Qinghai-Tibetan Plateau. *Ecology and Evolution* 3:4584-4595. doi: doi:10.1002/ece3.847
- Zuur A, Ieno EN, Walker N, Saveliev AA, Smith GM (2009) *Mixed effects models and extensions in ecology with R*. Springer, Berlin

## Tables

Table 1. The effect of water table depth and the most significant additional soil property (in terms of model fit) on the local species richness and phylogenetic clustering of the herbaceous and woody layer in savannas in southeast Brazil, using the generalized linear models. Models assumed negative binomial and Gaussian distributions, respectively.\* - $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$

Model	Fixed effects	Estimate	SE	<i>t</i>
Species Richness				
Herbaceous layer	log(water table depth)	0.20	0.06	3.65***
	soil P	-0.10	0.05	-1.90
Woody layer	log(water table depth)	2.09	0.30	6.97***
	soil P	-0.38	0.19	1.98*
Lineage Diversity				
Herbaceous layer	log(water table depth)	0.83	0.21	3.97***
	base saturation	0.37	0.21	1.78
Woody layer	log(water table depth)	-0.26	0.19	-1.39

Table 2. Results of the generalized additive models estimating the effect of groundwater depth on the local species richness and phylogenetic clustering of the herbaceous and woody layer in savannas in southeast Brazil. Models estimating effects on richness and lineage diversity assumed negative binomial distribution and Gaussian distributions, respectively. edf - effective degrees of freedom. \* - $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$

Model	edf	Chi.sq/F	R <sup>2</sup> adj
Richness			
Herbaceous layer	1.00	5.14*	0.24
Woody layer	3.55	75.8***	0.88
Lineage diversity			
Herbaceous layer	5.32	13.61***	0.82
Woody layer	1.00	0.00	-0.11

## Figure legends

**Fig 1.** Study site and experimental design. a) Satellite image of the Itirapina Ecological Station showing each of the 19 study sites, including marshy grasslands (rhombuses), wet grasslands (crosses) and savannas with low (squares), intermediate (triangles) and high (circles) woody cover. b) Design of a study site with the six 10 x 10 m sample plots (woody layer sampling) and detail of a plot with six the 1 x 1 m subplots (herbaceous layer sampling).

**Fig 2.** Relationship between species richness and lineage diversity of the herbaceous a,c) and woody layer b,d) and the water table depth in a Neotropical savanna in southeast Brazil. Negative values of lineage diversity indicate phylogenetic clustering and positive values indicate phylogenetic overdispersion. Lines are predicted values based on the results of generalized additive models and dashed lines indicate 95% confidence intervals

**Fig 3.** Occurrence of a) herbaceous and b) woody species along a water table depth gradient in Southeast Brazil, highlighting the distinct pattern of changes in species composition shown by herbaceous (turnover) and woody plant communities (nestedness). Species of each layer are ranked by their average water table occurrence ((minimum water table depth occurrence + maximum water table depth occurrence) / 2).

Fig 1.




Fig 2.




Fig 3.

